

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44, 31/535, C07D 213/38, 413/12	A1	(11) International Publication Number: WO 97/04778 (43) International Publication Date: 13 February 1997 (13.02.97)
(21) International Application Number: PCT/US96/11958 (22) International Filing Date: 18 July 1996 (18.07.96) (30) Priority Data: 60/001,588 27 July 1995 (27.07.95) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BRYANT, Henry, U. [US/US]; 7840 Camelback Drive, Indianapolis, IN 46250 (US). FINLEY, Don, R. [US/US]; 4644 Macy Drive, Greenwood, IN 46142 (US). MATSUMOTO, Ken [US/US]; 4240 Briarwood Drive, Indianapolis, IN 46250 (US). (74) Agents: STRODE, Janelle, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: PYRIDINE COMPOUNDS, INTERMEDIATES, PROCESSES, COMPOSITIONS, AND METHODS**(57) Abstract**

The present invention is directed to compounds of formula (I) wherein n is 2 or 3; R is dimethylamino, diethylamino, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, or 1-hexamethyleneimino; R¹ is hydrogen, loweralkyl of C₁-C₄, phenyl, or mono- or di-substituted phenyl wherein each substituent is independently halo, methyl, hydroxy, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₃-alkoxycarbonyloxy or C₄-C₆-alkylsulfonyloxy; R² is hydrogen, hydroxy, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₃-alkoxycarbonyloxy, or C₄-C₆-alkylsulfonyloxy; and the pharmaceutically acceptable acid addition salts, which are for the treatment of post menopausal symptoms such as osteoporosis, cardiovascular conditions such as hyperlipidaemia, and the like.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

PYRIDINE COMPOUNDS, INTERMEDIATES, PROCESSES, COMPOSITIONS,
AND METHODS

This invention relates to the fields of
5 pharmaceutical and organic chemistry and provides novel
pyridine compounds which are useful for the treatment of the
various medical indications associated with post-menopausal
syndrome. The present invention further relates to
intermediate compounds and processes useful for preparing the
10 pharmaceutically active compounds of the present invention,
and pharmaceutical compositions.

"Post-menopausal syndrome" is a term used to
describe various pathological conditions which frequently
15 affect women who have entered into or completed the
physiological metamorphosis known as menopause. Although
numerous pathologies are contemplated by the use of this
term, three major effects of post-menopausal syndrome are the
source of the greatest long-term medical concern:
20 osteoporosis, cardiovascular effects such as hyperlipidaemia,
and estrogen-dependent cancer, particularly breast and
uterine cancer.

Osteoporosis describes a group of diseases which
25 arise from diverse etiologies, but which are characterized by
the net loss of bone mass per unit volume. The consequence
of this loss of bone mass and resulting bone fracture is the
failure of the skeleton to provide adequate structural
support for the body. One of the most common types of
30 osteoporosis is that associated with menopause. Most women
lose from about 20% to about 60% of the bone mass in the
trabecular compartment of the bone within 3 to 6 years after
the cessation of menses. This rapid loss is generally
associated with an increase of bone resorption and formation.

-2-

However, the resorptive cycle is more dominant and the result is a net loss of bone mass. Osteoporosis is a common and serious disease among post-menopausal women.

There are an estimated 25 million women in the United States, alone, who are afflicted with this disease. The results of osteoporosis are personally harmful and also account for a large economic loss due its chronicity and the need for extensive and long term support (hospitalization and nursing home care) from the disease sequelae. This is especially true in more elderly patients. Additionally, although osteoporosis is not generally thought of as a life threatening condition, a 20% to 30% mortality rate is related with hip fractures in elderly women. A large percentage of this mortality rate can be directly associated with post-menopausal osteoporosis.

The most vulnerable tissue in the bone to the effects of post-menopausal osteoporosis is the trabecular bone. This tissue is often referred to as spongy or cancellous bone and is particularly concentrated near the ends of the bone (near the joints) and in the vertebrae of the spine. The trabecular tissue is characterized by small osteoid structures which inter-connect with each other, as well as the more solid and dense cortical tissue which makes up the outer surface and central shaft of the bone. This inter-connected network of trabeculae gives lateral support to the outer cortical structure and is critical to the bio-mechanical strength of the overall structure. In post-menopausal osteoporosis, it is, primarily, the net resorption and loss of the trabeculae which leads to the failure and fracture of bone. In light of the loss of the trabeculae in post-menopausal women, it is not surprising that the most common fractures are those associated with bones which are highly dependent on trabecular support, e.g., the vertebrae, the neck of the weight bearing bones such as the femur and the fore-arm. Indeed, hip fracture, collies fractures, and vertebral crush fractures are hall-marks of post-menopausal osteoporosis.

-3-

At this time, the only generally accepted method for treatment of post-menopausal osteoporosis is estrogen replacement therapy. Although therapy is generally successful, patient compliance with the therapy is low primarily because estrogen treatment frequently produces undesirable side effects.

Throughout premenopausal time, most women have less incidence of cardiovascular disease than age-matched men. Following menopause, however, the rate of cardiovascular disease in women slowly increases to match the rate seen in men. This loss of protection has been linked to the loss of estrogen and, in particular, to the loss of estrogen's ability to regulate the levels of serum lipids. The nature of estrogen's ability to regulate serum lipids is not well understood, but evidence to date indicates that estrogen can up regulate the low density lipid (LDL) receptors in the liver to remove excess cholesterol. Additionally, estrogen appears to have some effect on the biosynthesis of cholesterol, and other beneficial effects on cardiovascular health.

It has been reported in the literature that post-menopausal women having estrogen replacement therapy have a return of serum lipid levels to concentrations to those of the pre-menopausal state. Thus, estrogen would appear to be a reasonable treatment for this condition. However, the side-effects of estrogen replacement therapy are not acceptable to many women, thus limiting the use of this therapy. An ideal therapy for this condition would be an agent which would regulate the serum lipid level as does estrogen, but would be devoid of the side-effects and risks associated with estrogen therapy.

The third major pathology associated with post-menopausal syndrome is estrogen-dependent breast cancer and, to a lesser extent, estrogen-dependent cancers of other organs, particularly the uterus. Although such neoplasms are not solely limited to a post-menopausal women, they are more prevalent in the older, post-menopausal population. Current

-4-

chemotherapy of these cancers has relied heavily on the use of anti-estrogen compounds such as, for example, tamoxifen. Although such mixed agonist-antagonists have beneficial effects in the treatment of these cancers, and the estrogenic side-effects are tolerable in acute life-threatening situations, they are not ideal. For example, these agents may have stimulatory effects on certain cancer cell populations in the uterus due to their estrogenic (agonist) properties and they may, therefore, be contraproductive in some cases. A better therapy for the treatment of these cancers would be an agent which is an anti-estrogen compound having negligible or no estrogen agonist properties on reproductive tissues.

In response to the clear need for new pharmaceutical agents which are capable of alleviating the symptoms of, *inter alia*, post-menopausal syndrome, the present invention provides new pyridine compounds, pharmaceutical compositions thereof, and methods of using such compounds for the treatment of post-menopausal syndrome and other estrogen-related pathological conditions.

Smooth aortal muscle cell proliferation plays an important role in diseases such as atherosclerosis and restenosis. Vascular restenosis after percutaneous transluminal coronary angioplasty (PTCA) has been shown to be a tissue response characterized by an early and late phase. The early phase occurring hours to days after PTCA is due to thrombosis with some vasospasms while the late phase appears to be dominated by excessive proliferation and migration of aortal smooth muscle cells. In this disease, the increased cell motility and colonization by such muscle cells and macrophages contribute significantly to the pathogenesis of the disease. The excessive proliferation and migration of vascular aortal smooth muscle cells may be the primary mechanism to the reocclusion of coronary arteries following PTCA, atherectomy, laser angioplasty and arterial bypass graft surgery. See "Intimal Proliferation of Smooth Muscle Cells as an Explanation for Recurrent Coronary Artery

-5-

Stenosis after Percutaneous Transluminal Coronary Angioplasty," Austin et al., Journal of the American College of Cardiology, 8: 369-375 (Aug. 1985).


Vascular restenosis remains a major long term complication following surgical intervention of blocked arteries by percutaneous transluminal coronary angioplasty (PTCA), atherectomy, laser angioplasty and arterial bypass graft surgery. In about 35% of the patients who undergo PTCA, reocclusion occurs within three to six months after the procedure. The current strategies for treating vascular restenosis include mechanical intervention by devices such as stents or pharmacologic therapies including heparin, low molecular weight heparin, coumarin, aspirin, fish oil, calcium antagonist, steroids, and prostacyclin. These strategies have failed to curb the reocclusion rate and have been ineffective for the treatment and prevention of vascular restenosis. See "Prevention of Restenosis after Percutaneous Transluminal Coronary Angioplasty: The Search for a 'Magic Bullet'," Hermans et al., American Heart Journal, 122: 171-187 (July 1991).

In the pathogenesis of restenosis excessive cell proliferation and migration occurs as a result of growth factors produced by cellular constituents in the blood and the damaged arterial vessel wall which mediate the proliferation of smooth muscle cells in vascular restenosis.

Agents that inhibit the proliferation and/or migration of smooth aortal muscle cells are useful in the treatment and prevention of restenosis. The present invention provides for the use of compounds as smooth aortal muscle cell proliferation inhibitors and, thus inhibitors of restenosis.

-6-

The present invention is directed to compounds of Formula I:



wherein:

5 n is 2 or 3;

 R is dimethylamino, diethylamino, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, or 1-hexamethyleneimino;

 R¹ is hydrogen, loweralkyl of C₁-C₄, phenyl, or mono or di-substituted phenyl wherein each substituent is independently halo, methyl, hydroxy, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or C₄-C₆-alkylsulfonyloxy;

15 R² is hydrogen, hydroxy, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or C₄-C₆-alkylsulfonyloxy;

20 and the pharmaceutically acceptable acid addition salts thereof.

 These compounds are useful for the treatment of post menopausal symptoms such as osteoporosis, cardiovascular conditions such as hyperlipidaemia, and the like.

25 Accordingly, the present invention is also directed to methods employing and compositions comprising these compounds, for the treatment of post menopausal syndrome. The methods may additionally employ, and the compositions additionally comprise, an estrogen or progestin.

30 Preferred compounds are those with one or more of the following values, in any combination:

-7-

$n = 2$

$R = 1\text{-pyrrolidinyl}$


$R = 1\text{-piperidinyl}$

$R^1 = \text{methyl}$

5 $R^2 = \text{hydroxy}$

A particularly preferred compound is 2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-methylpyridine, including pharmaceutically acceptable salts.

Also provided by the present invention is a process for
10 preparing a compound of formula Ia:



wherein:


15 n is 2 or 3;

R is dimethylamino, diethylamino, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, or 1-hexamethyleneimino;

R^{1a} is hydrogen, loweralkyl of C₁-C₄, phenyl, or mono- or disubstituted phenyl wherein each substituent is
20 independently halo, methyl, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or C₄-C₆-alkylsulfonyloxy;

25 R^{2a} is hydrogen, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or C₄-C₆-alkylsulfonyloxy, which comprises reacting a compound
30 of Formula II:


-8-



II

5 with an alkyl chloride of the formula $\text{Cl}-(\text{CH}_2)_n\text{-R}$, in an inert solvent, in the presence of base, and at a temperature of from 0 to 100°C . The compounds of Formula II are another embodiment of the present invention.

10 The compounds of the present invention are prepared by the following alkylation reaction:



15 In Formula Ia and II, R^{1a} and R^{2a} have the same meaning as R^1 and R^2 , respectively, except that any hydroxy must be protected; if it is desired to obtain an R^2 =hydroxy compound, or an R^1 =hydroxy-substituted phenyl compound, the hydroxy group must first be protected, the alkylation reaction conducted, and the protection later removed.

20 Hydroxy protection is well known; see, for example, Protective Groups in Organic Chemistry, Plenum Press (London and New York, 1973); Protecting Groups in Organic Synthesis, Wiley (New York, 1981); and The Peptides, Vol. I, Schrooder and Lubke, Academic Press (London and New York, 1965).

Hydroxyl protection is also discussed in U.S. 4,418,068, which is incorporated herein by reference.

Suitable hydroxy protected groups are C₁-C₆ alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted
5 benzoyloxy as previously defined, C₁-C₅-alkoxycarbonyloxy, and C₄-C₆-alkylsulfonyloxy.

The alkylation reaction is carried out by contacting the reactants, typically in an inert solvent. Suitable such solvents include methyl ethyl ketone, acetone,
10 THF, dimethylformamide, and DMSO. A base is supplied to serve as a hydrogen halide acceptor. The reaction goes forward at temperatures over a wide range but is generally conducted at temperatures from 0 to 100°C. The reaction consumes the reactants and base in equimolar amounts;
15 preferably, the reactants are supplied in equimolar amounts and the base is supplied in an excess amount. Isolation and preparation are carried out in standard procedures.

Deprotection of the hydroxy protecting groups is carried out in known procedures, which are discussed in the
20 references cited above.

The compounds of Formula II are prepared according to the following scheme:

25 formula which is illustrated by preparations 1-16. In the



R^{1a} is hydrogen or loweralkyl and X is vinyl, for the
30 preparation of those compounds wherein R^{1a}=hydrogen or loweralkyl; or R^{1a} is phenyl or substituted phenyl and X is 2-chloroethyl, for the preparation of those compounds wherein R^{1a}=phenyl or substituted phenyl.

Although the free-base form of formula I compounds
35 can be used in the methods of the present invention, it is often preferred to prepare and use a pharmaceutically acceptable salt form. Thus, the compounds used in the

-10-

methods of this invention primarily form pharmaceutically acceptable acid addition salts with a wide variety of organic and inorganic acids, and include the physiologically acceptable salts which are often used in pharmaceutical chemistry. Such salts are also part of this invention. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, phosphoric, hypophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl substituted alkanoic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate, β -hydroxybutyrate, butyne-1,4-dioate, hexyne-1,4-dioate, caprate, caprylate, chloride, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, terephthalate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, sulfate, bisulfate, pyrosulfate, sulfite, bisulfite, sulfonate, benzenesulfonate, p-bromophenylsulfonate, chlorobenzenesulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, methanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like. A preferred salt is the hydrochloride salt.

The pharmaceutically acceptable acid addition salts are typically formed by reacting a compound of formula I with an equimolar or excess amount of acid. The reactants are generally combined in a mutual solvent such as diethyl ether or ethyl acetate. The salt normally precipitates out of

-11-

solution within about one hour to 10 days and can be isolated by filtration or the solvent can be stripped off by conventional means.

5 The pharmaceutically acceptable salts generally have enhanced solubility characteristics compared to the compound from which they are derived, and thus are often more amenable to formulation as liquids or emulsions.


The following examples are presented to further illustrate the preparation of compounds of the present
10 invention. It is not intended that the invention be limited in scope by reason of any of the following examples.

NMR data for the following Examples were generated on a GE 300 MHz NMR instrument, and anhydrous d-6 DMSO was used as the solvent unless otherwise indicated.

15

Preparation 1

2-(4-Benzoyloxyphenyl)-2-(N,N-diethylamino)acetonitrile



20


Diethylamine hydrochloride (76.7 g, 70 mmol) was added portion wise at rt to a 200 mL aqueous solution of potassium cyanide (45.5 g, 70 mmol) with stirring and warmed to 60-65 °C. A MeOH/THF (300 ml, 1:1) solution of 4-(benzyloxy)-
25 benzaldehyde (100 g, 47 mmol) (Aldrich) was added in a thin stream to the aqueous solution and stirred for 16 hr at ~62 °C. The reaction mixture was allowed to cool and the organics were removed in vacuo. The aqueous portion was diluted with an equal volume of brine and extracted several
30 times with EtOAc. The EtOAc extracts were combined, washed with H₂O, dried (Na₂SO₄), and concentrated to a tacky amber solid. The material was triturated with petroleum ether and

-12-

filtered to give a yellow solid, which was taken up in warm Et₂O, charcoaled, filtered, diluted with petroleum ether, and allowed to crystallize. The crystals were filtered and dried in vacuo to give 81.5 g (59%) of the desired product: mp 75-76 °C; ¹H NMR (CDCl₃) δ 1.06-1.11 (t, 6H, J = 7 Hz, CH₃), 2.42-2.49 (m, 2H, NCH₂), 2.66-2.73 (m, 2H, NCH₂), 4.98 (s, 1H, CHCN), 5.09 (s, 2H, OCH₂), 6.99-7.01 (d, 2H, J = 7 Hz, ArH), 7.35-7.49 (m, 7H, ArH); MS(FD) 294(M⁺). Anal. Calcd for C₁₉H₂₂N₂O: C, 77.52; H, 7.53; N, 9.51. Found: C, 77.70; H, 7.64; N, 9.45.

Preparation 2


4-Methoxybenzylchloride



15

4-Methoxybenzylalcohol (50 g, 0.36 mol) (Aldrich) was added dropwise to 100 mL of conc. HCl and stirred vigorously for 2 h at rt. The reaction mixture was extracted four times with Et₂O, the Et₂O extracts were combined, washed with brine, dried (Na₂SO₄), and concentrated. The yellow oil was distilled at 70-73 °C (0.2 mmHg) to give 52 g (93%) as a colorless oil: ¹H NMR (CDCl₃) δ 3.85 (s, 3H, OCH₃), 4.62 (s, 2H, CH₂Cl), 6.91-6.93 (d, 2H, J = 8 Hz, ArH), 7.38-7.40 (d, 2H, J = 8 Hz, ArH); MS(FD) 156(M⁺).

30


Preparation 31-(4-Benzoyloxyphenyl)-2-(4-methoxyphenyl)ethan-1-one

5

A 60% dispersion of sodium hydride in mineral oil (17 g, 0.42 mol) was washed with hexane and suspended in 200 mL DMF. 2-(4-benzoyloxyphenyl)-2-(N,N-diethylamino)acetonitrile (94 g, 0.32 mol) in 350 mL DMF was added dropwise over 1 h at rt to the NaH suspension and stirred an additional 1 h. A 300 mL DMF solution of 4-methoxybenzylchloride (50 g, 0.32 mol) was added dropwise over 45 min to the reaction mixture causing a mild exotherm to 45 °C. The reaction mixture was stirred for 16 h at rt, cooled in an ice bath, quenched with 100 mL MeOH, and reduced in vacuo to a smaller volume. The brown suspension was cooled in an ice bath and stirred while 2 L of 5 N HCl was added at a rate to keep the temperature below 25 °C. The suspension was stirred 18 h at rt, then extracted several times with CH₂Cl₂. The CH₂Cl₂ extracts were combined, washed with brine, dried (Na₂SO₄), and concentrated to give an oily brown solid. The product was triturated with Et₂O, filtered, and dried in vacuo to give 90 g (85%) of an off-white solid: mp 147-148 °C; ¹H NMR (CDCl₃) δ 3.80 (s, 3H, OCH₃), 4.18 (s, 2H, O=CCH₂), 5.13 (s, 2H, OCH₂Ar), 7.00-7.03 (d, 2H, J = 9 Hz, ArH), 7.18-7.21 (d, 2H, J = 9 Hz, ArH), 7.41-7.44 (m, 5H, ArH), 7.99-8.02 (d, 2H, J = 9 Hz, ArH); MS(FD) 332(M⁺). Anal. Calcd for C₂₂H₂₀O₃: C, 79.49; H, 6.06. Found: C, 79.23; H, 6.08.

30

-14-

Preparation 4**1-(4-Benzyloxyphenyl)-1-(dimethylamino)-2-(4-methoxyphenyl)ethylene**


5

Dimethylamine (55 mL, 0.15 mol) in 100 mL of toluene was added over ten min at 0-5 °C to a 600 mL toluene suspension of 1-(4-benzyloxyphenyl)-2-(4-methoxyphenyl)ethan-1-one (50 g, 0.15 mol), then stirred 15 min at 0 °C. A 50 mL toluene solution of TiCl_4 (9.8 mL, 0.09 mol) was added dropwise over 1 h to the reaction mixture, keeping the temperature below 5 °C. The brown suspension was stirred 16 h at rt then filtered through Hyflo Super Cel to give a yellow filtrate.

Two Et_2O washes of the filter bed were combined with the filtrate and the solution was concentrated to a thick amber oil. The oil was triturated with Et_2O / pet. ether and filtered to give 41 g (77%) of yellow solid: mp 60 °C; ^1H NMR (CDCl_3) d 2.64 (s, 6H, NMe_2), 3.71 (s, 3H, OCH_3), 5.07 (s, 2H, OCH_2), 5.42 (s, 1H, $=\text{CH}$), 6.57-6.60 (d, 2H, $J = 9$ Hz, ArH), 6.67-6.70 (d, 2H, $J = 9$ Hz, ArH), 6.86-6.89 (d, 2H, $J = 9$ Hz, ArH), 6.90-6.93 (d, 2H, $J = 9$ Hz, ArH), 7.18-7.21 (d, 2H, $J = 9$ Hz, ArH), 7.40-7.45 (m, 5H, ArH); MS(FD) $360[\text{M}+1]^+$. Anal. Calcd for $\text{C}_{24}\text{H}_{25}\text{NO}_2$: C, 80.19; H, 7.01; N, 3.90. Found: C, 80.41; H, 7.12; N, 4.09.

30

-15-

Preparation 5**1-(4-Benzoyloxyphenyl)-2-(4-methoxyphenyl)hexan-1,5-dione**

5

Methyl vinyl ketone (25 mL, 31 mmol) (Aldrich) was added to a 250 mL MeCN solution of 1-(4-benzyloxyphenyl)-1-dimethylamino-2-(4-methoxyphenyl)ethylene (28 g, 80 mmol) and refluxed 24 h. The reaction mixture was allowed to cool and concentrated to a gummy solid, which was then stirred for 16 h in 200 mL of 4N HCl. The aqueous mixture was extracted several times with EtOAc. The EtOAc extracts were combined, washed with brine, and concentrated to a solid. Recrystallisation from EtOAc/ hexane gave 27.3 g (87%) of the desired product: mp 93-94 °C; ¹H NMR (CDCl₃) δ 2.08-2.41 (comp, 7H, O=CCH₃, CH₂CH₂), 3.75 (s, 3H, OCH₃), 4.54-4.57 (t, 1H, J = 4, Hz, CH), 5.08 (s, 2H, OCH₂Ar), 6.80-6.83 (d, 2H, J = 9 Hz, ArH), 6.91-6.94 (d, 2H, J = 9 Hz, ArH), 7.16-7.19 (d, 2H, J = 9 Hz, ArH), 7.34-7.39 (m, 5H, ArH), 7.92-7.95 (d, 2H, J = 9 Hz, ArH); MS(FD) 402(M⁺). Anal. Calcd for C₂₆H₂₆O₄: C, 77.79; H, 6.61. Found: C, 77.59; H, 6.51.

Preparation 6**1-(4-Benzoyloxyphenyl)-2-(4-methoxyphenyl)-5-phenylpentan-1,5-dione**

25

1-(4-Benzoyloxyphenyl)-1-dimethylamino-2-(4-methoxyphenyl)ethylene (5.03 g, 14 mmol) and 3-chloropropiophenone (2.47 g, 15 mmol) (Aldrich) were refluxed in 60 mL MeCN for 18 h, then worked up and purified to give 5.18 g (79%) of product: mp 76-78 °C; ¹H NMR (CDCl₃) δ 2.22-

30


-16-

3.00 (comp, 4H, CH₂CH₂), 3.75 (s, 3H, OCH₃), 4.64-4.69 (t, 1H, *J* = 7 Hz, CH), 5.08 (s, 2H, OCH₂Ar), 6.80-6.83 (d, 2H, *J* = 9 Hz, ArH), 6.92-6.94 (d, 2H, *J* = 9 Hz, ArH), 7.20-7.54 (comp, 10H, ArH), 7.89-7.98 (m, 4H, ArH); MS(FD) 464(M⁺).
 5 Anal. Calcd for C₃₁H₂₈O₄: C, 80.15; H, 6.07. Found: C, 80.38; H, 6.15.

Preparation 7

1-(4-Benzyloxyphenyl)-2-(4-methoxyphenyl)-5-(4-fluorophenyl)pentan-1,5-dione

10




The 1-(4-benzyloxyphenyl)-1-dimethylamino-2-(4-methoxyphenyl)ethylene (5.4 g, 15 mmol) and 3-chloro-4'-fluoropropiophenone (3.1 g, 17 mmol) (Aldrich) were refluxed in 60 mL MeCN for 20 h and then the reaction mixture worked up and purified to give 4.2 g (66%) of white solid: mp 90-93 °C; ¹H NMR (CDCl₃) δ 2.21-2.98 (comp, 4H, CH₂CH₂), 3.76 (s, 3H, OCH₃), 4.63-4.68 (t, 1H, *J* = 7 Hz, CH), 5.09 (s, 2H, OCH₂Ar), 6.81-6.84 (d, 2H, *J* = 9 Hz, ArH), 6.92-6.95 (d, 2H, *J* = 9 Hz, ArH), 7.08-7.40 (comp, 9H, ArH), 7.92-7.98 (m, 4H, ArH); FD(MS) 482(M⁺). Anal. Calcd for C₃₁H₂₇O₄F: C, 77.16; H, 5.64. Found: C, 77.22; H, 5.72.

25

30

-17-

Preparation 8**1-(4-Benzyloxyphenyl)-2,5-bis(4-methoxyphenyl)pentan-1,5-dione**


5

The 1-(4-benzyloxyphenyl)-1-dimethylamino-2-(4-methoxyphenyl)ethylene (10 g, 28 mmol) and 3-chloro-4'-methoxypropiophenone (6.5 g, 33 mmol) were refluxed in 150 mL MeCN for 24 h, then the reaction mixture was worked up and purified to give 10 g (72%) of product: mp 132-134 °C; ¹H NMR (CDCl₃) δ 2.20-2.98 (comp, 4H, CH₂CH₂), 3.75 (s, 3H, OCH₃), 4.64-4.69 (t, 1H, J = 7 Hz, CH), 5.08 (s, 2H, OCH₂Ar), 6.81-6.94 (m, 6H, ArH), 7.17-7.42 (m, 7H, ArH), 7.87-8.00 (m, 4H, ArH); MS(FD) 495[M+1]⁺. Anal. Calcd for C₃₂H₃₀O₅: C, 77.71; H, 6.11. Found: C, 77.80; H, 6.05.

15

Preparation 9**2-(4-Benzyloxyphenyl)-3-(4-methoxyphenyl)-6-methylpyridine**

20




1-(4-Benzyloxyphenyl)-2-(4-methoxyphenyl)hexan-1,5-dione (10 g, 0.025 mol) was dissolved in 300 mL MeOH,

-18-

hydroxylamine hydrochloride (2.16 g, 0.031 mol) was added, and the reaction mixture was refluxed for 18 h. The reaction mixture was allowed to cool, most of the MeOH removed in vacuo, and then was diluted with 250 mL water. The aqueous mixture was extracted several times with EtOAc, the EtOAc extracts were combined, washed with brine, dried (Na₂SO₄), and concentrated. The residue was purified by prep. chromatography (EtOAc/ hexane gradient) and recrystallised from EtOAc/ hexane to give 4.4 g (47%) of product: mp 135-136 °C; ¹H NMR (CDCl₃) δ 2.74 (s, 3H, Pyr-CH₃), 3.92 (s, 3H, OCH₃), 5.16 (s, 2H, OCH₂Ar), 6.91-6.98 (m, 4H, ArH), 7.18-7.25 (m, 3H, ArH), 7.37-7.52 (m, 7H, ArH), 7.66-7.68 (d, 1H, J = 8 Hz, ArH); MS(FD) 381(M⁺). Anal. Calcd for C₂₆H₂₃NO₂: C, 81.86; H, 6.08; N, 3.67. Found: C, 82.10, H, 6.21; N, 3.71.

Preparation 10

2-(4-Benzyloxyphenyl)-3-(4-methoxyphenyl)-6-phenylpyridine




The 1-(benzyloxyphenyl)-2-(4-methoxyphenyl)-5-phenylpentan-1,5-dione (1.0 g, 2.2 mmol) and hydroxylamine hydrochloride (0.3 g, 4.3 mmol) were refluxed in 30 mL of MeOH for 48 h, worked up, purified, and recrystallised from Et₂O/ pet. ether to give 0.32 g (34 % yield) of white solid: mp 130-132 °C; ¹H NMR (CDCl₃) δ 3.82 (s, 3H, OCH₃), 5.06 (s, 2H, OCH₂Ar), 6.84-6.90 (m, 4H, ArH), 7.15-7.18 (d, 2H, J = 9, ArH), 7.38-7.49 (comp, 10H, ArH), 7.73 (s, 2H, ArH), 8.12-8.15 (d, 2H, J = 9, ArH); MS(FD) 443(M⁺). Anal. Calcd for

-19-

C₃₁H₂₅NO₂: C, 83.94; H, 5.68; N, 3.16. Found: C, 84.14; H, 5.90; N, 2.94.

Preparation 11

5 **2-(4-Benzyloxyphenyl)-3-(4-methoxyphenyl)-6-(4-fluorophenyl)pyridine**




10 The 1-(4-benzyloxyphenyl)-2-(4-methoxyphenyl)-5-(4-fluorophenyl)pentan-1,5-dione (1.0 g, 2.4 mmol) and hydroxylamine hydrochloride (0.2 g, 2.9 mmol) were refluxed in 30 mL of MeOH for 48 h, worked up, and purified to give
 15 0.45 g (40 % yield) of product: mp 114-115 °C; ¹H NMR (CDCl₃) δ 3.82 (s, 3H, OCH₃), 5.06 (s, 2H, OCH₂Ar), 6.83-6.89 (m, 4H, ArH), 7.13-7.19 (m, 4H, ArH), 7.34-7.44 (m, 7H, ArH), 7.65-7.74 (dd, 2H, J = 8, ArH), 8.10-8.14 (m, 2H, ArH); MS(FD) 461(M⁺). Anal. Calcd for C₃₁H₂₄NO₂F: C, 80.67; H, 5.24; N, 3.03. Found: C, 80.87; H, 5.30; N, 3.09.

20

Preparation 12

2-(4-Benzyloxyphenyl)-3,6-bis(4-methoxyphenyl)pyridine




-20-

The 1-(4-benzyloxyphenyl)-2,5-bis(4-methoxyphenyl)pentan-1,5-dione (9 g, 18 mmol), hydroxylamine hydrochloride (1.8 g, 26 mmol), and 0.8 mL concd H₂SO₄ were stirred in 250 mL of 1-butanol at 95-100 °C for 18 h. The reaction mixture was neutralized with solid NaHCO₃, worked up, purified, and recrystallised (EtOAc/ hexane) to give 3.2 g (37% yield) of white solid: mp 99-102 °C; ¹H NMR (CDCl₃) δ 3.82 (s, 3H, OCH₃), 3.87 (s, 3H, OCH₃), 5.06 (s, 2H, OCH₂Ar), 6.83-6.89 (m, 4H, ArH), 6.99-7.02 (d, 2H, J = 9, ArH), 7.14-7.17 (d, 2H, J = 9, ArH), 7.32-7.45 (m, 7H, ArH), 7.64-7.71 (m, 2H, ArH), 8.08-8.11 (d, 2H, J = 9, ArH); MS(FD) 473 (M⁺). Anal. Calcd for C₃₂H₂₇NO₃: C, 81.16; H, 5.75; N, 2.96. Found: C, 81.10; H, 5.92; N, 2.98.

Preparation 13

2-(4-Hydroxyphenyl)-3-(4-methoxyphenyl)-6-methylpyridine



20


The 2-(4-benzyloxyphenyl)-3-(4-methoxyphenyl)-6-methylpyridine (14 g, .37 mmol) and ammonium formate (3.5 g, 55 mmol) were stirred in 1 L of EtOH, then 5% Pd/ C (5.8 g, 3 mmol Pd) was added and the reaction mixture warmed to 50-55 °C. The reaction mixture was stirred at 55 °C for 1 h at which time the TLC (MeOH/ CHCl₃, 1:9) indicated the starting material was consumed. After cooling, the reaction mixture was filtered, concentrated, and the residue triturated with Et₂O to give 9.6 g (90% yield) of white solid: mp 190 °C; ¹H NMR (CDCl₃) δ 2.66 (s, 3H, Pyr-CH₃), 3.82 (s, 3H, OCH₃), 6.53-6.56 (d, 2H, J = 8 Hz, ArH), 6.80-6.83 (d, 2H, J = 8

30

-21-

Hz, ArH), 7.06-7.18 (m, 6H, ArH, ArOH), 7.59-7.62 (d, 1H, J = 8 Hz, ArH); MS(FD) 291(M^+). Anal. Calcd for $C_{19}H_{17}NO_2$: C, 78.33; H, 5.88; N, 4.81. Found: C, 78.11; H, 5.93; N, 4.72.

5


Preparation 142-(4-Hydroxyphenyl)-3-(4-methoxyphenyl)-6-phenylpyridine

10

The 2-(4-benzyloxyphenyl)-3-(4-methoxyphenyl)-6-phenylpyridine (3.5 g, 7.9 mmol) and ammonium formate (2.5 g, 40 mmol) were stirred in 150 mL of EtOH, then 5% Pd/C (1.4 g, 0.6 mmol Pd) was added and the reaction mixture was stirred at 60 °C for 1h. The TLC (MeOH/ $CHCl_3$, 1:9) indicated a small amount of the starting material remained, so more ammonium formate (0.3 g, 5 mmol) was added and reaction mixture stirred at 60 °C for 0.5 h, which consumed the remaining starting material. After cooling, the reaction mixture was filtered and the filter pad washed with 50 mL of EtOH, then 50 mL of EtOAc. The filtrates were combined and concentrated, then redissolved in 150 mL of EtOAc, washed with brine, dried (Na_2SO_4), concentrated, and the residue triturated with Et₂O/ pet. ether to give 2.5 g (90% yield) of white solid: mp 200-202 °C; 1H NMR ($CDCl_3$) d 3.85 (s, 3H, OCH₃), 5.07 (s, 1H, ArOH), 6.71-6.90 (m, 4H, ArH), 7.17-7.20 (d, 2H, J = 8 Hz, ArH), 7.38-7.54 (m, 5H, ArH), 7.75 (s, 2H, ArH), 8.13-8.16 (d, 2H, J = 9 Hz, ArH); MS(FD) 354(M^+). Anal. Calcd for $C_{24}H_{19}NO_2$: C, 81.56; H, 5.42; N, 3.96. Found: C, 81.50; H, 5.61; N, 3.99.


Preparation 15**2-(4-Hydroxyphenyl)-3-(4-methoxyphenyl)-6-(4-fluorophenyl)pyridine**

5



The 2-(4-benzyloxyphenyl)-3-(4-methoxyphenyl)-6-(4-fluorophenyl)pyridine (1.2 g, 2.6 mmol) was dissolved in 50 mL of THF/ EtOH (1:1), 5% Pd/ C (0.3 g, 0.14 mmol Pd) added, and the reaction mixture hydrogenated at 40 psi for 4 h at rt. The reaction mixture was filtered and concentrated to give 0.86 g (90% yield) of white foam: mp 143-145 °C; ¹H NMR (CDCl₃) δ 3.83 (s, 3H, OCH₃), 4.78 (s, 1H, ArOH), 6.72-6.75 (d, 2H, *J* = 8 Hz, ArH), 6.84-6.87 (d, 2H, *J* = 8 Hz, ArH), 7.14-7.27 (m, 4H, ArH), 7.37-7.40 (d, 2H, *J* = 8 Hz, ArH), 7.69-7.73 (m, 2H, ArH), 8.10-8.15 (m, 2H, ArH); MS(FD) 371(M⁺). Anal. Calcd for C₂₄H₁₈NO₂F: C, 77.61; H, 4.89; N, 3.77. Found: C, 77.88; H, 5.12; N, 3.51.

20


Preparation 16**2-(4-Hydroxyphenyl)-3,6-bis(4-methoxyphenyl)pyridine**

-23-

The 2-(4-benzyloxyphenyl)-3,6-bis(4-methoxyphenyl)pyridine (3.0 g, 6.3 mmol), ammonium formate (2.0 g, 31 mmol), and 5% Pd/C (1.0 g, 0.5 mmol Pd) were combined in 150 mL of EtOH, and the reaction mixture was stirred at 50-55 °C and followed by TLC (EtOAc/ Tol, 1:9). After 1 h, more ammonium formate (0.5 g, 8 mmol) was added and reaction mixture stirred at 50 °C for 30 min and no starting material remained. After cooling, the reaction mixture was filtered, concentrated, and the residue trituated with Et₂O/ pet. ether. The white ppt was filtered and dried in vacuo to give 2.35g (97% yield) of product: mp 204-205 °C; ¹H NMR (CDCl₃) δ 3.83 (s, 3H, OCH₃), 3.88 (s, 3H, OCH₃), 4.85 (s, 1H, ArOH), 6.71-6.74 (d, 2H, J = 9 Hz, ArH), 6.83-6.86 (d, 2H, J = 9 Hz, ArH), 7.00-7.03 (d, 2H, J = 9 Hz, ArH), 7.14-7.17 (d, 2H, J = 9 Hz, ArH), 7.37-7.40 (d, 2H, J = 9 Hz, ArH), 7.68-7.70 (m, 2H, ArH), 8.08-8.11 (d, 2H, J = 9 Hz, ArH); MS(FD) 383(M⁺). Anal. Calcd for C₂₅H₂₁NO₃: C, 78.31; H, 5.52; N, 3.65. Found: C, 78.07; H, 5.79; N, 3.95.

Example 1

2-[4-[2-(1-N,N-Dimethylamino)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine




The 2-(4-hydroxyphenyl)-3-(4-methoxyphenyl)-6-methylpyridine (3, 5g, 12 mmol), 2-dimethylaminoethylchloride hydrochloride (1.73 g, 12 mmol), and powdered K₂CO₃ (8.3 g, 60 mmol) were stirred in 250 mL methyl ethyl ketone and

30

-24-

refluxed 20 h. The reaction mixture was worked up and purified by prep. chromatography (EtOH/ CH₂Cl₂ gradient) to give 3.7 g (85% yield) as a clear gum. A portion of the gum (1.6 g, 4.4 mmol) was dissolved in 50 mL MeOH, 30 mL HCl/ MeOH was added at 0 °C, stirred 10 min, worked up and triturated with Et₂O, then filtered and dried in vacuo to give 1.48 g (84 % conversion of free base to salt) as the HCl salt monohydrate: mp 135-140 °C; ¹H NMR (CDCl₃) δ 1.60-1.90 (bs, 2H, H₂O), 2.92-2.93 (bs, 6H, NCH₃), 3.13 (s, 3H, pyr-CH₃), 3.47-3.49 (m, 2H, CH₂N), 3.82 (s, 3H, ArOCH₃), 4.50-4.53 (m, 2H, OCH₂), 6.85-6.90 (m, 4H, ArH), 7.00-7.04 (m, 2H, ArH), 7.41-7.44 (m, 2H, ArH), 7.54-7.56 (d, 1H, J = 8 Hz, ArH), 8.12-8.14 (d, 1H, J = 8 Hz, ArH); MS(FAB) 363 (M⁺). Anal. Calcd for C₂₃H₂₆N₂O₂ · HCl · H₂O: C, 66.26; H, 7.01; N, 6.72. Found: C, 66.54; H, 6.72; N, 6.79.

Example 2**2-[4-[2-(1-Pyrrolidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine**


The 2-(4-hydroxyphenyl)-3-(4-methoxyphenyl)-6-methylpyridine (4 g, 14 mmol), N-(2-chloroethyl)pyrrolidine hydrochloride (2.3 g, 14 mmol), and powdered K₂CO₃ (9.5 g, 68 mmol) were stirred in 250 mL methyl ethyl ketone and refluxed 18 h. The reaction mixture was worked up and purified by prep. chromatography (EtOH/ CH₂Cl₂ gradient), triturated with Et₂O, filtered, and dried in vacuo to give 3.7 g (71% yield) of white product: mp 63-64 °C; ¹H NMR (CDCl₃) δ

-25-

1.79-1.83 (comp, 4H, pyrrolidine), 2.60-2.64 (comp, 7H, Pyr-CH₃ and pyrrolidine), 2.86-2.90 (t, 2H, J = 6 Hz, NCH₂), 3.81 (s, 3H, ArOCH₃), 4.07-4.11 (t, 2H, J = 6 Hz, OCH₂), 6.78-6.82 (m, 4H, ArH), 7.06-7.30 (comp, 5H, ArH), 7.55-7.58 (d, 1H, J = 8 Hz, ArH); MS(FD) 388(M⁺). Anal. Calcd for C₂₅H₂₈N₂O₂: C, 77.23; H, 7.26; N, 7.21. Found: C, 77.12; H, 7.48; N, 7.00.

Example 3

10 2-[4-[2-(4-Morpholinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine




15

The 2-(4-hydroxyphenyl)-3-(4-methoxyphenyl)-6-methylpyridine (3.0 g, 10 mmol), N-(2-chloroethyl)morpholine hydrochloride (1.9 g, 10 mmol), and powdered K₂CO₃ (7.1 g, 51 mmol) were stirred in 200 mL methyl ethyl ketone and refluxed 20 h. The reaction mixture was worked up and purified by prep. chromatography (EtOH/ CH₂Cl₂ gradient), triturated with Et₂O/ pet. ether, filtered, and dried in vacuo to give 3.4 g (82% yield) of white solid: mp 66-67 °C; ¹H NMR (CDCl₃) δ 2.55-2.58 (m 4H, morpholine), 2.62 (s, 3H, pyr-CH₃), 2.76-2.80 (t, 2H, J = 6 Hz, NCH₂), 3.72-3.75 (m, 4H, morpholine), 3.80 (s, 3H, ArOCH₃), 4.06-4.10 (t, 2H, J = 6 Hz, OCH₂), 6.75-6.81 (m, 4H, ArH), 7.05-7.14 (m, 4H, ArH), 7.26-7.29 (m, 1H, ArH) 7.54-7.56 (d, 1H, J = 8 Hz, ArH); MS(FD) 404(M⁺). Anal. Calcd for C₂₅H₂₈N₂O₃: C, 74.23; H, 6.98; N, 6.93. Found: C, 74.50; H, 6.91; N, 7.05.

-26-

Example 4**2-[4-[2-(1-Pyrrolidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-phenylpyridine**

5




The 2-(4-hydroxyphenyl)-3-(4-methoxyphenyl)-6-phenylpyridine (2.0 g, 5 mmol), N-(2-chloroethyl)pyrrolidine hydrochloride (1.0 g, 5 mmol), and powdered K_2CO_3 (4.2 g, 30 mmol) were stirred in 200 mL methyl ethyl ketone and refluxed 20 h. The reaction mixture was worked up and the residue triturated with pet. ether, filtered, and dried in vacuo to give 2.1 g (84% yield) of an off-white solid: mp 100-103.
15 °C; 1H NMR ($CDCl_3$) δ 1.81-1.84 (m, 4H, pyrrolidine), 2.61-2.65 (m, 4H, pyrrolidine), 2.88-2.92 (t, 2H, $J = 6$ Hz, NCH_2), 3.83 (s, 3H, $ArOCH_3$), 4.10-4.14 (t, 2H, $J = 6$ Hz, OCH_2), 6.81-6.87 (m, 4H, ArH), 7.15-7.18 (d, 2H, $J = 9$ Hz, ArH) 7.42-7.50 (m, 5H, ArH), 7.73 (s, 2H, ArH), 8.13-8.16 (d, 2H, $J = 9$ Hz, ArH); MS(FD) 450(M^+). Anal. Calcd for $C_{30}H_{30}N_2O_2$: C, 79.97; H, 6.71; N, 6.22. Found: C, 79.86; H, 6.09; N, 6.16.
20

25

Example 5**2-[4-[2-(1-Pyrrolidinyl)ethoxy]phenyl]-3,6-bis(4-methoxyphenyl)pyridine**

5




The 2-(4-hydroxyphenyl)-3,6-bis(4-methoxyphenyl)pyridine (2.3 g, 6 mmol), N-(2-chloroethyl)pyrrolidine hydrochloride (1.1 g, 6 mmol), and powdered K_2CO_3 (4.1 g, 30 mmol) were stirred in 250 mL methyl ethyl ketone and refluxed 18 h. The TLC (EtOH/ CH_2Cl_2 , 1:4) of the reaction mixture indicated some starting material remained, so more N-(2-chloroethyl)pyrrolidine hydrochloride (0.3 g, 1.6 mmol) and K_2CO_3 (2 g, 15 mmol) were added and reaction mixture refluxed an additional 18 h. The reaction mixture was worked up, concentrated, and the residue was dissolved in 25 mL MeOH, treated with 5 mL HCl/ MeOH at 0 °C, and concentrated. The residue was recrystallised from MeOH/ Et_2O to give 1.8 g (58 % yield) of product as the HCl salt: mp 230-231 °C; 1H NMR ($CDCl_3$) δ 1.61 (bs, 2H, pyrrolidine), 2.10-2.26 (comp, 4H, pyrrolidine), 2.90-3.10 (bs, 2H, pyrrolidine), 3.48 (bs, 2H, NCH_2), 3.84 (s, 3H, $ArOCH_3$), 3.88 (s, 3H, $ArOCH_3$), 4.10-4.14 (m, 2H, OCH_2), 6.78-6.81 (d, 2H, J = 9 Hz, ArH), 6.86-6.89 (d, 2H, J = 9 Hz, ArH), 7.00-7.03 (d, 2H, J = 9 Hz, ArH), 7.14-7.16 (d, 2H, J = 9 Hz, ArH), 7.44-7.46 (d, 2H, J = 9 Hz, ArH), 7.66-7.73 (m, 2H, ArH), 8.07-8.10 (d, 2H, J = 9 Hz, ArH); MS(FD) 480(M^+). Anal. Calcd for $C_{31}H_{32}N_2O_3 \cdot HCl$:

-28-

C, 72.01; H, 6.43; N, 5.42. Found: C, 72.02; H, 6.65; N, 5.24.

Example 6

5 2-[4-[2-(1-Piperidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-(4-fluorophenyl)pyridine




10 The 2-(4-hydroxyphenyl)-3-(4-methoxyphenyl)-6-(4-fluorophenyl)pyridine (1.78 g, 4.8 mmol), N-(2-chloroethyl)piperidine hydrochloride (0.88 g, 4.8 mmol), and powdered K_2CO_3 (3.3 g, 24 mmol) were stirred in 150 mL methyl ethyl ketone and refluxed 18 h. The reaction mixture was

15 worked up and concentrated to a light yellow gum. The gum was warmed in 50 mL of acetone, citric acid·H₂O (1 equivalent) was added, the clear solution warmed 5 min, then concentrated, triturated with Et₂O, filtered, and dried in vacuo to give 1.1 g (48% overall yield) of product as the

20 citrate salt: mp 100 °C; ¹H NMR (CDCl₃) δ 1.25-1.52 (comp, 6H, piperidine), 2.38-2.54 (m, 4H, citrate), 2.73-2.74 (m, 4H, piperidine), 2.2.93-2.95 (bs, 2H, CH₂N), 3.49 (s, 3H, ArOCH₃), 3.97-4.00 (bs, 2H, OCH₂), 6.47-6.53 (m, 4H, ArH), 6.80-6.87 (m, 4H, ArH), 7.07-7.10 (m, 2H, ArH), 7.40-7.41 (m, 2H, ArH), 7.78-7.82 (m, 2H, ArH); MS(FAB) 483 [M+]⁺ Anal. Calcd for C₃₁H₃₁N₂O₂F · C₆H₈O₇: C, 65.87; H, 5.83; N, 4.15.

25 Found: C, 65.90; H, 6.12; N, 4.09.

Example 7**2-[4-[2-(1-Piperidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine**

Sodium hydride (60 % in mineral oil, 0.26 g, 6.6 mmol) was washed with 2 x 5 mL hexane, then added portionwise to a solution of the 2-(4-hydroxyphenyl)-3-(4-methoxyphenyl)-6-methylpyridine (1.75 g, 6 mmol) in 60 mL of DMF and stirred at 60 °C for 1 h. A 30 mL DMF solution of N-(2-chloroethyl)piperidine (1.06 g, 7.2 mmol) was added dropwise to the reaction mixture and stirred at 55-60 °C for 8 h, followed by cooling and quenching with 250 mL of ice water.

15 The aqueous mixture was extracted with EtOAc (3 x 100 mL) and 100 mL Et₂O and the organics were combined, washed with brine, dried (Na₂SO₄), then concentrated to an oil. The oil was taken up 100 mL Et₂O, treated with decolorizing carbon, filtered, and concentrated to give 2.1 g (87 % yield) of product as a gum. A portion of the gum (1.3 g, 3.3 mmol) was dissolved in acetone (100 mL), citric acid monohydrate (0.7 g, 3.3 mmol) added, and mixture warmed at gentle reflux for 10 min, then concentrated. The residue was triturated with Et₂O, filtered, and dried in vacuo to give 1.7 g (90 % conversion of free base to salt) as the solid citrate salt: mp 101-110 °C; ¹H NMR (CDCl₃ + DMSO-d₆) δ 1.56-1.85 (comp, 6H, piperidine), 2.56 (s, 3H, pyr-CH₃), 2.68-2.84 (m, 4H, citrate), 3.06-3.12 (m, 4H, piperidine), 3.28-3.30 (m, 2H, CH₂N), 3.74 (s, 3H, ArOCH₃), 4.27-4.30 (m, 2H, OCH₂), 6.69-6.77 (m, 4H, ArH), 6.99-7.25 (comp, 5H, ArH), 7.50-7.53 (d,

20

25

30


-30-

^1H , $J = 8 \text{ Hz}$, ArH); MS(FAB) 403 $[\text{M}+1]^+$. Anal. Calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_2 \cdot \text{C}_6\text{H}_8\text{O}_7$: C, 64.63; H, 6.44; N, 4.71. Found: C, 64.39; H, 6.44; N, 4.93.

5

Example 8


2-[4-[2-(N,N-Dimethylamino)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-methylpyridine



10


The AlCl_3 (7 g, 52 mmol) was stirred in 150 mL of DCE at 0 °C, ethanethiol (5.5 mL, 75 mmol) was added, and the mixture stirred for 15 min. The 2-[4-[2-(N,N-dimethylamino)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine (2.7 g, 7.5 mmol) in 50 mL DCE was added dropwise to the reaction mixture and stirred for 2 h as it was allowed to come to rt, then quenched with 25 mL THF at 0 °C, followed by 25 mL 1 N HCl and worked up. The crude product was recrystallised from hot EtOAc/ pet. ether to give 0.9 g (34 % yield) of white solid: mp 155-157 °C; ^1H NMR (CDCl_3) δ 2.37 (s, 6H, NCH_3), 2.62 (s, 3H, pyr-CH_3), 2.78-2.81 (t, 2H, $J = 5 \text{ Hz}$, CH_2N), 4.05-4.07 (t, 2H, $J = 5 \text{ Hz}$, OCH_2), 6.64-6.69 (m, 4H, ArH), 6.94-6.96 (d, 2H, $J = 8 \text{ Hz}$, ArH), 7.09-7.12 (d, 1H, $J = 8 \text{ Hz}$, ArH), 7.24-7.27 (comp, 2H, ArH), 7.52-7.55 (d, 1H, $J = 8 \text{ Hz}$, ArH); MS(FD) 348 (M^+). Anal. Calcd for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_2$: C, 75.84; H, 6.94; N, 8.04. Found: C, 75.70; H, 6.98; N, 8.00.

-31-

Example 9**2-[4-[2-(1-Pyrrolidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-methylpyridine**

The AlCl_3 (3.6 g, 27 mmol) was stirred in 100 mL of DCE at 0 °C, ethanethiol (3 mL, 38 mmol) was added, and the mixture stirred for 15 min. The 2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine (1.5 g, 3.9 mmol) in 50 mL DCE was added dropwise to the reaction mixture and stirred for 2 h as it was allowed to come to rt, then quenched with 75 mL THF at 0 °C, followed by 50 mL 1 N HCl and worked up. The crude product was triturated with Et_2O , filtered, and dried in vacuo to give 1.3 g (90 % yield) of an off-white solid: mp 159-161 °C; ^1H NMR (CDCl_3) δ 1.80-1.84 (bs, 4H, pyrrolidine), 2.61 (s, 3H, pyr- CH_3), 2.67-2.69 (bs, 4H, pyrrolidine), 2.93-2.97 (m, 2H, CH_2N), 4.08-4.10 (m, 2H, OCH_2), 6.60-6.65 (m, 4H, ArH), 6.90-6.93 (d, 2H, $J = 8$ Hz, ArH), 7.08-7.11 (d, 1H, $J = 8$ Hz, ArH), 7.23-7.26 (m, 2H, ArH), 7.50-7.53 (d, 1H, $J = 8$ Hz, ArH); MS(FD) 374 (M^+). Anal. Calcd for $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_2$: C, 76.98; H, 7.00; N, 7.48. Found: C, 76.70; H, 7.07; N, 7.36.

-32-

Example 10**2-[4-[2-(1-Piperidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-methylpyridine**

The AlCl_3 (1.2 g, 9 mmol) was stirred in 30 mL of DCE at 0 °C, ethanethiol (1 mL, 14 mmol) was added, and the mixture stirred for 15 min. The 2-[4-[2-(1-

10 piperidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine (0.5 g, 1.2 mmol) in 20 mL DCE was added dropwise to the reaction mixture and stirred for 2 h as it was allowed to come to rt, then quenched with 25 mL THF at 0 °C, followed by 25 mL 1 N HCl and worked up. The free base


15 gave a very clean NMR, but the elemental analysis was low for carbon and hydrogen, so it was taken up in 50 mL hot acetone, citric acid \cdot H_2O (0.2 g, 0.9 mmol) was added, then heated 10 min, concentrated, triturated with Et_2O , filtered, and dried in vacuo to give 0.38 g (72 % yield) of product as the

20 citrate salt: mp 90-100 °C; ^1H NMR (CDCl_3) (free base) δ 1.34-1.66 (comp, 6H, piperidine), 2.42-2.65 (comp, 7H, pyr- CH_3 and piperidine), 2.70-2.79 (m, 2H, CH_2N), 3.95-4.10 (m, 2H, OCH_2), 6.50-6.59 (m, 4H, ArH), 6.76-6.85 (d, 2H, J = 8 Hz, ArH), 6.98-7.05 (d, 1H, J = 8 Hz, ArH), 7.10-7.21 (m, 2H, ArH), 7.41-7.50 (d, 1H, J = 8 Hz, ArH); MS(FAB) 398 $[\text{M}+1]^+$. Anal. Calcd for $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_2 \cdot \text{C}_6\text{H}_8\text{O}_7$: C, 64.13; H, 6.25; N, 4.82. Found: C, 63.99; H, 6.36; N, 4.58.

-33-

Example 11

2-[4-[2-(4-Morpholinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-methylpyridine




The AlCl_3 (5 g, 38 mmol) was stirred in 100 mL of DCE at 0 °C, ethanethiol (4 mL, 54 mmol) was added, and the mixture stirred for 15 min. The 2-[4-[2-(4-morpholinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine (2.2 g, 5.4 mmol) in 50 mL DCE was added dropwise to the reaction mixture and stirred for 2 h as it was allowed to come to rt, then quenched with 50 mL THF at 0 °C, followed by 50 mL 1 N HCl and worked up. The crude product was triturated with Et_2O / pet. ether, filtered, and dried in vacuo to give 1.9 g (90 % yield) of white solid: mp 75-80 °C; ^1H NMR (CDCl_3) δ 2.59-2.62 (comp, 7H, morpholine and pyr- CH_3), 2.79-2.83 (t, 2H, J = 5 Hz, CH_2N), 3.74-3.77 (m, 4H, morpholine), 4.07-4.11 (t, 2H, J = 5 Hz, OCH_2), 6.63-6.71 (m, 4H, ArH), 6.94-6.97 (d, 2H, J = 8 Hz, ArH), 7.11-7.13 (d, 1H, J = 8 Hz, ArH), 7.24-7.27 (d, 2H, J = 8 Hz, ArH), 7.53-7.56 (d, 1H, J = 8 Hz, ArH); MS(FD) 390 (M^+). Anal. Calcd for $\text{C}_{24}\text{H}_{26}\text{N}_3\text{O}_3$: C, 73.82; H, 6.71; N, 7.17. Found: C, 73.75; H, 6.71; N, 7.05.

-34-

Example 12

2-[4-[2-(1-Piperidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-(4-fluorophenyl)pyridine




The AlCl_3 (1.9 g, 14 mmol) was stirred in 50 mL of DCE at 0 °C, ethanethiol (1.5 mL, 20 mmol) was added, and the mixture stirred for 15 min. The 2-[4-[2-(1-piperidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-(4-fluorophenyl)pyridine (1 g, 2 mmol) in 25 mL DCE was added dropwise to the reaction mixture and stirred for 2 h as it was allowed to come to rt, then quenched with 25 mL THF at 0 °C, followed by 25 mL 1 N HCl and worked up. The crude product was triturated with Et_2O , filtered, and dried in vacuo to give 0.86 g (91 % yield) of white solid: mp 194-195 °C; ^1H NMR (CDCl_3) δ 1.63-1.67 (comp, 6H, piperidine), 2.59 (bs, 4H, piperidine), 2.81-2.84 (t, 2H, J = 5 Hz, CH_2N), 4.12-4.15 (t, 2H, J = 5 Hz, OCH_2), 6.64-6.70 (m, 4H, ArH), 7.00-7.03 (d, 2H, J = 8 Hz, ArH), 7.12-7.18 (m, 2H, ArH), 7.37-7.40 (d, 2H, J = 8 Hz, ArH), 7.61-7.67 (m, 2H, ArH), 8.08-8.12 (m, 2H, ArH); MS (FD) 468 (M^+). Anal. Calcd for $\text{C}_{30}\text{H}_{29}\text{N}_2\text{O}_2$ F: C, 76.90; H, 6.24; N, 5.98. Found: C, 77.13; H, 6.36; N, 5.86.

25

-35-

Example 13

2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-phenylpyridine



5


The 2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-phenylpyridine as the HCl salt (1.9 g, 3.9 mmol) was slurried with 50 mL of DCE in a screw cap flask, cooled to 0 °C, and stirred while the BCl₃ (2.7 mL at 0 °C, 32 mmol) in 5 mL DCE was added and the reaction flask sealed. After 2 h, TLC (EtOH/ CH₂Cl₂, 1:4) indicated that starting material remained, so an additional 1.7 mL of BCl₃ (20 mmol) was added and reaction mixture stirred for 18 h. The reaction mixture was carefully quenched with 25 mL MeOH at 0 °C, poured in 150 mL of aqueous NaHCO₃, and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ extracts were combined, washed with brine, dried (Na₂SO₄), concentrated, and recrystallised from EtOH to give 1.35 g (79 % yield) of product: mp 176-178 °C; ¹H NMR (CDCl₃) δ 1.86 (bs, 4H, pyrrolidine), 2.73 (bs, 4H, pyrrolidine), 2.97-3.01 (t, 2H, J = 5 Hz, CH₂N), 4.12-4.16 (t, 2H, J = 5 Hz, OCH₂), 6.64-6.67 (m, 4H, ArH), 6.98-7.01 (d, 2H, J = 8 Hz, ArH), 7.38-7.50 (m, 5H, ArH), 7.67 (s, 2H, ArH), 8.10-8.12 (d, 2H, J = 8 Hz, ArH); MS(FD) 436 (M⁺). Anal. Calcd for C₂₉H₂₈N₂O₂: C, 79.79; H, 6.46; N, 6.42. Found: C, 80.01; H, 6.55; N, 6.54.

10

15

20

25

Example 14**2-[4-[2-(1-Pyrrolidinyl)ethoxy]phenyl]-3,6-bis(4-hydroxyphenyl)pyridine**

The 2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3,6-bis(4-methoxyphenyl)pyridine as the HCl salt (1.1 g, 2.1 mmol) was
 10 slurried with 50 mL of DCE in a screw cap flask, cooled to 0 °C, and stirred while the BCl₃ (5.5 mL at 0 °C, 64 mmol) was added, the reaction flask sealed, and reaction mixture stirred for 20 h. The reaction mixture was carefully quenched with 25 mL MeOH at 0 °C, poured in 150 mL of aqueous
 15 NaHCO₃, and extracted with CH₂Cl₂ (3 x 100 mL). The CH₂Cl₂ extracts were combined, washed with brine, dried (Na₂SO₄), concentrated, and recrystallised from EtOH to give 0.89g (83 % yield) of product: mp 145-150 °C; ¹H NMR (CDCl₃) δ 1.63 (bs, 2H, pyrrolidine), 2.01-2.76 (comp, 6H, pyrrolidine and
 20 CH₂N), 3.90-3.94 (m, 2H, OCH₂), 6.55-6.60 (m, 4H, ArH), 6.6.71-6.84 (m, 4H, ArH), 7.19-7.22 (d, 2H, J = 9 Hz, ArH) 7.43-7.45 (m, 2H, ArH), 7.76-7.79 (d, 2H, J = 9 Hz, ArH); MS(FD) 452 (M⁺). Anal. Calcd for C₂₉H₂₈N₂O₃: C, 76.97; H, 6.24; N, 6.19. Found: C, 76.92; H, 6.18; N, 6.13.

-37-

In the examples illustrating the methods, a post-menopausal model was used in which effects of different treatments upon circulating lipids were determined.

Seventy-five day old female Sprague Dawley rats (weight range of 200 to 225g) were obtained from Charles River Laboratories (Portage, MI). The animals were either bilaterally ovariectomized (OVX) or exposed to a sham surgical procedure at Charles River Laboratories, and then shipped after one week. Upon arrival, they were housed in metal hanging cages in groups of 3 or 4 per cage and had ad libitum access to food (calcium content approximately 0.5%) and water for one week. Room temperature was maintained at $22.2^{\circ} \pm 1.7^{\circ} \text{C}$ with a minimum relative humidity of 40%. The photoperiod in the room was 12 hours light and 12 hours dark.

Dosing Regimen Tissue Collection. After a one week acclimation period (therefore, two weeks post-OVX) daily dosing with test compound was initiated. 17α -ethynyl estradiol or the test compound were given orally, unless otherwise stated, as a suspension in 1% carboxymethylcellulose or dissolved in 20% cyclodextrin. Animals were dosed daily for 4 days. Following the dosing regimen, animals were weighed and anesthetized with a ketamine: Xylazine (2:1, V:V) mixture and a blood sample was collected by cardiac puncture. The animals were then sacrificed by asphyxiation with CO_2 , the uterus was removed through a midline incision, and a wet uterine weight was determined.

Cholesterol Analysis. Blood samples were allowed to clot at room temperature for 2 hours, and serum was obtained following centrifugation for 10 minutes at 3000 rpm. Serum cholesterol was determined using a Boehringer Mannheim Diagnostics high performance cholesterol assay. Briefly the

-38-

cholesterol was oxidized to cholest-4-en-3-one and hydrogen peroxide. The hydrogen peroxide was then reacted with phenol and 4-aminophenazone in the presence of peroxidase to produce a p-quinone imine dye, which was read spectrophotometrically at 500 nm. Cholesterol concentration was then calculated against a standard curve. The entire assay was automated using a Biomek Automated Workstation.

Uterine Eosinophil Peroxidase (EPO) Assay. Uteri were kept at 4° C until time of enzymatic analysis. The uteri were then homogenized in 50 volumes of 50 mM Tris buffer (pH - 8.0) containing 0.005% Triton X-100. Upon addition of 0.01% hydrogen peroxide and 10 mM O-phenylenediamine (final concentrations) in Tris buffer, increase in absorbance was monitored for one minute at 450 nm. The presence of eosinophils in the uterus is an indication of estrogenic activity of a compound. The maximal velocity of a 15 second interval was determined over the initial, linear portion of the reaction curve.

Source of Compound: 17 α -ethynyl estradiol was obtained from Sigma Chemical Co., St. Louis, MO.

Influence of Formula I Compounds on Serum Cholesterol and Determination of Agonist/Non-Agonist Activity

Data presented in Table 1 below show comparative results among ovariectomized rats, rats treated with 17 α -ethynyl estradiol (EE₂; an orally available form of estrogen), and rats treated with certain compounds of the present invention. Although EE₂ caused a decrease in serum cholesterol when orally administered at 0.1 mg/kg/day, it also exerted a stimulatory action on the uterus so that EE₂ uterine weight was substantially greater than the uterine weight of ovariectomized control animals. This uterine response to estrogen is well recognized in the art.

Not only did the compounds of the present invention generally reduce serum cholesterol compared to the

-39-

ovariectomized control animals, but uterine weight was only minimally increased to slightly decreased with the majority of the formula compounds tested. Compared to estrogenic compounds known in the art, the benefit of serum cholesterol reduction without adversely affecting uterine weight is quite rare and desirable.

As is expressed in the below data, estrogenicity also was assessed by evaluating the response of eosinophil infiltration into the uterus. Most of the compounds of the present invention did not cause any increase in the number of eosinophils observed in the stromal layer of ovariectomized rats, while others caused only modest increases. Estradiol caused a substantial, expected increase in eosinophil infiltration.

The data presented in Table 1 below reflects the response of 5 to 6 rats per treatment. "NA" indicates that a compound was not active at the doses tested.

Table 1

Compound	Dose (mg/kg)	Uterine Wt. (%) inc. vs. OVX)	Uterine EPO (V. max)	Serum Cholesterol (% dec. vs. OVX)	ED50 (effective dose to dec. serum chol. by 50%)
EE2	0.1	221.8*	235.6*	88.3*	
Example 6	1	15.6	3.7	12.3	
	10	44.2	5.8	-9.1	
	25	69.5*	22.6*	40.8*	70
EE2	0.1	183.3*	339.2*	86.9*	
Example 6	0.1	9.5	6.6	4	
	1	-2.4	5.9	7.4	
	10	47.6*	6.5	33.1*	105

-40-

EE2	0.1	122.3*	79.0*	83.1*	
Example 12	0.1	9.8	5.9	26.2	
	1	4.2	9.7	9.8	
	10	76.2*	43.9*	0.1	NA
EE2	0.1	125.8*	44.9*	83.3*	
Example 12	1	2.5	5.4	24.7*	
	10	62.5*	70.1*	60.6*	
	25	68.0*	67.9*	49.9*	1.5
EE2	0.1	118.2*	153.9*	96.4*	
Example 5	0.1	-6.5	4.5	2.4	
	1	-7.6	4.2	14	
	10	27.2*	23.1	23.5*	>10
EE2	0.1	118.2*	153.9*	96.4*	
Example 14	0.1	6.3	2.4	11.9	
	1	-3.3	4.5	-16.5	
	10	8.8	4.2	-7.3	NA
EE2	0.1	105.6*	140.4*	71.4*	
Example 13	1	54.0*	61.8*	53.8*	
	10	38.8*	39.0	36.8*	
	25	35.5*	53.4*	62.6*	<1
EE2	0.1	93.8*	207.6*	68.0*	
Example 13	0.1	23.8	4.5	1.8	
	1	61.9*	34.8	50.9*	
	10	47.4*	32.1	55.6*	
	25	46.6*	22.2	39.4*	ED50~1
EE2	0.1	125.8*	111.2*	88.1	
Example 3	1	4.9	10.8*	4.5	
	10	-6.1	6	1.4	
	25	-2.3	3.8	-11.6	NA

-41-

EE2	0.1	144.0*	185.7*	90.1*	
Example 11	1	-0.8	4.8	16	
	10	27	4.9	18.2	
	25	5	4.4	-3.6	NA
EE2	0.1	104.5*	20.4*	90.5*	
Example 10	1	3.4	4.4	8.4	
	10	7	3.4	1.7	
	25	47.0*	6.8*	17.9	NA
EE2	0.1	82.2*	192.9	85.4*	
Example 7	0.1	-2.2	4.1	23.2	
	1	64.4*	19.1*	51.8*	
	10	-2.2	2.5	25.9	1
EE2	0.1	132.1*	349.8*	90.5*	
Example 7	0.1	0.4	7.3	27.7*	
	1	-0.4	17.3*	17.5	
	10	10.7	9.1	34.0*	11
EE2	0.1	144.0*	185.7*	90.1*	
Example 7	1	16.4	3.1	-35.5*	
	10	5.4	3.6	-5.2	
	25	-5.6	3.6	-1.2	NA
EE2	0.1	102.2*	62.7*	82.9*	
Example 2	1	12.7	6.6	-23.6	
	10	-12.5	5.9	-33.5	
	25	19.5*	59.8	9.7	NA
EE2	0.1	125.8*	44.9*	83.3*	
Example 9	1	2	6.7	23*	
	10	26.3*	10.2	49.2*	
	25	24.3*	8.5	58.5*	2.2

-42-

EE2	0.1	144.0*	185.7*	90.1*	
Example 9	1	6.2	4.6	-2	
	10	30.3*	4.4	29.5*	
	25	35.1*	5.6	50.0*	28
EE2	0.1	223.2*	265.2*	91.2*	
Example 1	1	0.6	2.9	15	
	10	2.6	2.5	14.9	
	25	5.2	2.3	22.2*	>1000
EE2	0.1	144.0*	185.7*	90.1*	
Example 8	1	-1.5	5	15.7	
	10	-1	6.7	1.5	
	25	41.6*	58.6*	46.4*	26

*P = ≤ 0.5 ANOVA with post-hoc Fisher's PLSD on raw data

5 In addition to the demonstrated benefits of the compounds of the present invention, especially when compared to estradiol, the above data clearly demonstrate that compounds of Formula I are not estrogen mimetics. Furthermore, no deleterious toxicological effects (survival) were observed with any treatment.

Osteoporosis Test Procedure

Following the General Preparation Procedure, *infra*, the rats were treated daily for 35 days (6 rats per treatment group) and sacrificed by carbon dioxide asphyxiation on the 36th day. The 35 day time period was sufficient to allow maximal reduction in bone density, measured as described herein. At the time of sacrifice, the uteri were removed, dissected free of extraneous tissue, and the fluid contents were expelled before determination of wet weight in order to confirm estrogen deficiency associated with complete ovariectomy. Uterine weight was routinely reduced about 75% in response to ovariectomy.

-43-

The right femurs were excised and digitized x-rays generated and analyzed by an image analysis program (NIH image) at the distal metaphysis. The proximal aspect of the tibiae from these animals were also scanned by quantitative computed tomography.

In accordance with the above procedures, the compound of Example 9 and ethynyl estradiol (EE₂) in 20% hydroxypropyl β -cyclodextrin were orally administered to test animals. Distal femur metaphysis data presented in Tables 2 and 3 below are the results of compound treatments compared to intact and ovariectomized test animals. The rats in the Table 2 tests were 6 months of age rather than 75 days of age (at the beginning of the tests).

15

Table 2

<u>Compound</u>	<u>Dose</u> (mg/kg)	<u>Body</u> Wt. Change (% dec. vs. OVX)	<u>Uterine</u> Wt. (% inc. vs. OVX)	<u>Femur</u> (x-ray image) (% protect.)	<u>Serum</u> Chol. (% dec. vs. OVX)
EE2	0.1	22.4*	128.3*	41.8*	73.8*
Example 9	1	8.3	-8.6	5.9	35.4*
	10	4.5	4.3	-15.3	35.4*
	25	6.3	-6.8	33.3	19.7
EE2	0.1	59.1*	226.1*	25.5*	73.6*
Example 9	0.01	-5	-3.5	-11.5	44.4*
	0.1	-8.2	-11	-16.4*	33.6*
	1	13.4	17.4	-9.9	54.2*
	10	29.6	62.0	28.1	42.2*

*P = \leq 0.5 ANOVA with post-hoc Fisher's PLSD on raw data

20

In summary, ovariectomy of the test animals caused a significant reduction in femur density compared to intact, vehicle treated controls. Orally administered ethynyl

-44-

estradiol (EE₂) prevented this loss, but the risk of uterine stimulation with this treatment is ever-present.

The compound of Example 9 of the present invention elevated mean bone density in a general, dose-related manner. Accordingly, the compounds of the present invention are useful for the treatment of post-menopausal syndrome, particularly osteoporosis.

MCF-7 Proliferation Assay

10 MCF-7 breast adenocarcinoma cells (ATCC HTB 22) were maintained in MEM (minimal essential medium, phenol red-free, Sigma, St. Louis, MO) supplemented with 10% fetal bovine serum (FBS) (V/V), L-glutamine (2 mM), sodium pyruvate (1 mM), HEPES ((N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid])10 mM), non-essential amino acids and 15 bovine insulin (1 ug/mL) (maintenance medium). Ten days prior to assay, MCF-7 cells were switched to maintenance medium supplemented with 10% dextran coated charcoal stripped fetal bovine serum (DCC-FBS) assay medium) in place of 10% 20 FBS to deplete internal stores of steroids. MCF-7 cells were removed from maintenance flasks using cell dissociation medium (Ca⁺⁺/Mg⁺⁺ free HBSS (phenol red-free) supplemented with 10 mM HEPES and 2 mM EDTA). Cells were washed twice with assay medium and adjusted to 80,000 cells/mL.

25 Approximately 100 μ L (8,000 cells) were added to flat-bottom microculture wells (Costar 3596) and incubated at 37° C in a 5% CO₂ humidified incubator for 48 hours to allow for cell adherence and equilibration after transfer. Serial dilutions of drugs or DMSO as a diluent control were prepared in assay 30 medium and 50 μ L transferred to triplicate microcultures followed by 50 μ L assay medium for a final volume of 200 μ L. After an additional 48 hours at 37° C in a 5% CO₂ humidified incubator, microcultures were pulsed with tritiated thymidine (1 uCi/well) for 4 hours. Cultures were terminated by 35 freezing at -70° C for 24 hours followed by thawing and harvesting of microcultures using a Skatron Semiautomatic Cell Harvester. Samples were counted by liquid scintillation

-45-

using a Wallac BetaPlace β counter. Results in Table 4 below show the IC₅₀ for certain compounds of the present invention.

Table 4

5

Compound (Example Reference)	IC ₅₀ nM
1	1000
2	1000
3	1000
5	1000
6	470
7	1000
8	1000
9	1000
11	1000
12	185
14	9

Uterine Fibrosis Test Procedures

Test 1

10 Between 3 and 20 women having uterine fibrosis are administered a compound of the present invention. The amount of compound administered is from 0.1 to 1000 mg/day, and the period of administration is 3 months.

15 The women are observed during the period of administration, and up to 3 months after discontinuance of administration, for effects on uterine fibrosis.

Test 2

20 The same procedure is used as in Test 1, except the period of administration is 6 months.

Test 3

 The same procedure is used as in Test 1, except the period of administration is 1 year.

-46-

Test 4

A. Induction of fibroid tumors in guinea pig.

Prolonged estrogen stimulation is used to induce
5 leiomyomata in sexually mature female guinea pigs. Animals
are dosed with estradiol 3-5 times per week by injection for
2-4 months or until tumors arise. Treatments consisting of a
compound of the invention or vehicle is administered daily
for 3-16 weeks and then animals are sacrificed and the uteri
10 harvested and analyzed for tumor regression.

B. Implantation of human uterine fibroid tissue in nude mice.

Tissue from human leiomyomas are implanted into the
15 peritoneal cavity and or uterine myometrium of sexually
mature, castrated, female, nude mice. Exogenous estrogen are
supplied to induce growth of the explanted tissue. In some
cases, the harvested tumor cells are cultured in vitro prior
to implantation. Treatment consisting of a compound of the
20 present invention or vehicle is supplied by gastric lavage on
a daily basis for 3-16 weeks and implants are removed and
measured for growth or regression. At the time of sacrifice,
the uteri are harvested to assess the status of the organ.

25 Test 5

A. Tissue from human uterine fibroid tumors is harvested
and maintained, in vitro, as primary nontransformed cultures.
Surgical specimens are pushed through a sterile mesh or
sieve, or alternately teased apart from surrounding tissue to
30 produce a single cell suspension. Cells are maintained in
media containing 10% serum and antibiotic. Rates of growth
in the presence and absence of estrogen are determined.
Cells are assayed for their ability to produce complement
component C3 and their response to growth factors and growth
35 hormone. In vitro cultures are assessed for their
proliferative response following treatment with progestins,
GnRH, a compound of the present invention and vehicle.

-47-

Levels of steroid hormone receptors are assessed weekly to determine whether important cell characteristics are maintained in vitro. Tissue from 5-25 patients are utilized.

- Activity in at least one of the above tests
- 5 indicates the compounds of the present invention are of potential in the treatment of uterine fibrosis.

Endometriosis Test Procedure

- In Tests 1 and 2, effects of 14-day and 21-day
- 10 administration of compounds of the present invention on the growth of explanted endometrial tissue can be examined.

Test 1

- Twelve to thirty adult CD strain female rats are
- 15 used as test animals. They are divided into three groups of equal numbers. The estrous cycle of all animals is monitored. On the day of proestrus, surgery is performed on each female. Females in each group have the left uterine horn removed, sectioned into small squares, and the squares
- 20 are loosely sutured at various sites adjacent to the mesenteric blood flow. In addition, females in Group 2 have the ovaries removed.

- On the day following surgery, animals in Groups 1 and 2 receive intraperitoneal injections of water for 14 days
- 25 whereas animals in Group 3 receive intraperitoneal injections of 1.0 mg of a compound of the present invention per kilogram of body weight for the same duration. Following 14 days of treatment, each female is sacrificed and the endometrial explants, adrenals, remaining uterus, and ovaries, where
- 30 applicable, are removed and prepared for histological examination. The ovaries and adrenals are weighed.

Test 2

- Twelve to thirty adult CD strain female rats are
- 35 used as test animals. They are divided into two equal groups. The estrous cycle of all animals is monitored. On the day of proestrus, surgery is performed on each female.

-48-

Females in each group have the left uterine horn removed, sectioned into small squares, and the squares are loosely sutured at various sites adjacent to the mesenteric blood flow.

- 5 Approximately 50 days following surgery, animals assigned to Group 1 receive intraperitoneal injections of water for 21 days whereas animals in Group 2 receive intraperitoneal injections of 1.0 mg of a compound of the present invention per kilogram of body weight for the same
10 duration. Following 21 days of treatment, each female is sacrificed and the endometrial explants and adrenals are removed and weighed. The explants are measured as an indication of growth. Estrous cycles are monitored.

15 Test 3

A. Surgical induction of endometriosis

- Autographs of endometrial tissue are used to induce endometriosis in rats and/or rabbits. Female animals at reproductive maturity undergo bilateral oophorectomy, and
20 estrogen is supplied exogenously thus providing a specific and constant level of hormone. Autologous endometrial tissue is implanted in the peritoneum of 5-150 animals and estrogen supplied to induce growth of the explanted tissue. Treatment consisting of a compound of the present invention is supplied
25 by gastric gavage on a daily basis for 3-16 weeks, and implants are removed and measured for growth or regression. At the time of sacrifice, the intact horn of the uterus is harvested to assess status of endometrium.

30 B. Implantation of human endometrial tissue in nude mice.

- Tissue from human endometrial lesions is implanted into the peritoneum of sexually mature, castrated, female, nude mice. Exogenous estrogen is supplied to induce growth of the explanted tissue. In some cases, the harvested
35 endometrial cells are cultured in vitro prior to implantation. Treatment consisting of a compound of the present invention supplied by gastric gavage on a daily basis

-49-

for 3-16 weeks, and implants are removed and measured for growth or regression. At the time of sacrifice, the uteri are harvested to assess the status of the intact endometrium.

5 Test 4

A. Tissue from human endometrial lesions is harvested and maintained *in vitro* as primary nontransformed cultures. Surgical specimens are pushed through a sterile mesh or sieve, or alternately teased apart from surrounding tissue to
10 produce a single cell suspension. Cells are maintained in media containing 10% serum and antibiotic. Rates of growth in the presence and absence of estrogen are determined. Cells are assayed for their ability to produce complement component C3 and their response to growth factors and growth
15 hormone. *In vitro* cultures are assessed for their proliferative response following treatment with progestins, GnRH, a compound of the invention, and vehicle. Levels of steroid hormone receptors are assessed weekly to determine whether important cell characteristics are maintained in
20 *vitro*. Tissue from 5-25 patients is utilized.

Inhibition of Aortal Smooth Cell Proliferation/Restenosis
Test Procedure

Compounds of the present invention have capacity to
25 inhibit aortal smooth cell proliferation. This can be demonstrated by using cultured smooth cells derived from rabbit aorta, proliferation being determined by the measurement of DNA synthesis. Cells are obtained by explant method as described in Ross, J. of Cell Bio. 50: 172 (1971).
30 Cells are plated in 96 well microtiter plates for five days. The cultures become confluent and growth arrested. The cells are then transferred to Dulbecco's Modified Eagle's Medium (DMEM) containing 0.5 - 2% platelet poor plasma, 2 mM L-glutamine, 100 U/ml penicillin, 100 mg/ml streptomycin, 1
35 mC/ml ³H-thymidine, 20 ng/ml platelet-derived growth factor, and varying concentrations of the present compounds. Stock solution of the compounds is prepared in dimethyl sulphoxide

-50-

and then diluted to appropriate concentration (0.01 - 30 mM) in the above assay medium. Cells are then incubated at 37° C. for 24 hours under 5% CO₂/95% air. At the end of 24 hours, the cells are fixed in methanol. ³H thymidine incorporation in DNA is then determined by scintillation counting as described in Bonin, et al., Exp. Cell Res. 181: 475-482 (1989).

Inhibition of aortal smooth muscle cell proliferation by the compounds of the present invention are further demonstrated by determining their effects on exponentially growing cells. Smooth muscle cells from rabbit aortae are seeded in 12 well tissue culture plates in DMEM containing 10% fetal bovine serum, 2 mM L-glutamine, 100 U/ml penicillin, and 100 mg/ml streptomycin. After 24 hours, the cells are attached and the medium is replaced with DMEM containing 10% serum, 2 mM L-glutamine, 100 U/ml penicillin, 100 mg/ml streptomycin, and desired concentrations of the compounds. Cells are allowed to grow for four days. Cells are treated with trypsin and the number of cells in each culture is determined by counting using a ZM-Coulter counter.

Activity in the above tests indicates that the compounds of the present invention are of potential in the treatment of restenosis.

The present invention also provides a method of alleviating post-menopausal syndrome in women which comprises the aforementioned method using compounds of Formula I and further comprises administering to a woman an effective amount of estrogen or progestin. These treatments are particularly useful for treating osteoporosis and lowering serum cholesterol because the patient will receive the benefits of each pharmaceutical agent while the compounds of the present invention would inhibit undesirable side-effects of estrogen and progestin. Activity of these combination treatments in any of the post-menopausal tests, infra, indicates that the combination treatments are useful for alleviating the symptoms of post-menopausal symptoms in women.

-51-

As used herein, the term "estrogen" includes steroidal compounds having estrogenic activity such as, for example, 17 β -estradiol, estrone, conjugated estrogen (Premarin®), equine estrogen, 17 α -ethynyl estradiol, and the like. As used herein, the term "progestin" includes compounds having progestational activity such as, for example, progesterone, norethynodrel, norgestrel, megestrol acetate, norethindrone, and the like.

Various forms of estrogen and progestin are commercially available. Estrogen-based agents include, for example, ethynyl estrogen (0.01 - 0.03 mg/day), mestranol (0.05 - 0.15 mg/day), and conjugated estrogenic hormones such as Premarin® (Wyeth-Ayerst; 0.3 - 2.5 mg/day). Progestin-based agents include, for example, medroxyprogesterone such as Provera® (Upjohn; 2.5 -10 mg/day), norethynodrel (1.0 - 10.0 mg/day), and norethindrone (0.5 - 2.0 mg/day). A preferred estrogen-based compound is Premarin, and norethynodrel and norethindrone are preferred progestin-based agents.

The method of administration of each estrogen- and progestin-based agent is consistent with that which is known in the art. For the majority of the methods of the present invention, compounds of Formula I are administered continuously, from 1 to 3 times daily. However, cyclical therapy may especially be useful in the treatment of endometriosis or may be used acutely during painful attacks of the disease. In the case of restenosis, therapy may be limited to short (1-6 months) intervals following medical procedures such as angioplasty.

As used herein, the term "effective amount" means an amount of compound of the present invention which is capable of alleviating the symptoms of the various pathological conditions herein described. The specific dose of a compound administered according to this invention will, of course, be determined by the particular circumstances surrounding the case including, for example, the compound administered, the route of administration, the state of being

-52-

of the patient, and the pathological condition being treated. A typical daily dose will contain a nontoxic dosage level of from about 5 mg to about 1000 mg/day of a compound of the present invention. Preferred daily doses generally will be
5 from about 15 mg to about 500 mg/day.

The compounds of this invention can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds preferably are formulated prior to
10 administration, the selection of which will be decided by the attending physician. Thus, another aspect of the present invention is a pharmaceutical composition comprising an effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, optionally
15 containing an effective amount of estrogen or progestin, and a pharmaceutically acceptable carrier, diluent, or excipient.

The total active ingredients in such formulations comprises from 0.1% to 99.9% by weight of the formulation. By "pharmaceutically acceptable" it is meant the carrier,
20 diluent, excipients and salt must be compatible with the other ingredients of the formulation, and not deleterious to the recipient thereof.

Pharmaceutical formulations of the present invention can be prepared by procedures known in the art
25 using well known and readily available ingredients. For example, the compounds of formula I, with or without an estrogen or progestin compound, can be formulated with common excipients, diluents, or carriers, and formed into tablets, capsules, suspensions, powders, and the like. Examples of
30 excipients, diluents, and carriers that are suitable for such formulations include the following: fillers and extenders such as starch, sugars, mannitol, and silicic derivatives; binding agents such as carboxymethyl cellulose and other cellulose derivatives, alginates, gelatin, and polyvinyl-
35 pyrrolidone; moisturizing agents such as glycerol; disintegrating agents such as calcium carbonate and sodium bicarbonate; agents for retarding dissolution such as

-53-

paraffin; resorption accelerators such as quaternary ammonium compounds; surface active agents such as cetyl alcohol, glycerol monostearate; adsorptive carriers such as kaolin and bentonite; and lubricants such as talc, calcium and magnesium stearate, and solid polyethyl glycols.

The compounds also can be formulated as elixirs or solutions for convenient oral administration or as solutions appropriate for parenteral administration, for example, by intramuscular, subcutaneous or intravenous routes.

10 Additionally, the compounds are well suited to formulation as sustained release dosage forms and the like. The formulations can be so constituted that they release the active ingredient only or preferably in a particular physiological location, possibly over a period of time. The

15 coatings, envelopes, and protective matrices may be made, for example, from polymeric substances or waxes.

Compounds of formula I, alone or in combination with a pharmaceutical agent of the present invention, generally will be administered in a convenient formulation.

20 The following formulation examples only are illustrative and are not intended to limit the scope of the present invention.

Formulations

In the formulations which follow, "active ingredient" means a compound of Formula I, or a salt or solvate thereof.

25

Formulation 1: Gelatin Capsules

30 Hard gelatin capsules are prepared using the following:

Ingredient	Quantity (mg/capsule)
Active ingredient	0.1 - 1000
Starch, NF	0 - 650
Starch flowable powder	0 - 650
Silicone fluid 350 centistokes	0 - 15

-54-

The formulation above may be changed in compliance with the reasonable variations provided.

A tablet formulation is prepared using the ingredients below:

5

Formulation 2: Tablets

Ingredient	Quantity (mg/tablet)
Active ingredient	2.5 - 1000
Cellulose, microcrystalline	200 - 650
Silicon dioxide, fumed	10 - 650
Stearate acid	5 - 15

10 The components are blended and compressed to form tablets.

Alternatively, tablets each containing 2.5 - 1000 mg of active ingredient are made up as follows:

Formulation 3: Tablets

15

Ingredient	Quantity (mg/tablet)
Active ingredient	25 - 1000
Starch	45
Cellulose, microcrystalline	35
Polyvinylpyrrolidone (as 10% solution in water)	4
Sodium carboxymethyl cellulose	4.5
Magnesium stearate	0.5
Talc	1

The active ingredient, starch, and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh
20 U.S. sieve. The granules so produced are dried at 50°-60° C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc,

-55-

previously passed through a No. 60 U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets.

Suspensions each containing 0.1 - 1000 mg of
5 medicament per 5 ml dose are made as follows:

Formulation 4: Suspensions

Ingredient	Quantity (mg/5 ml)
Active ingredient	0.1 - 1000 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mg
Benzoic acid solution	0.10 mL
Flavor	q.v.
Color	q.v.
Purified water to	5 mL

10 The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor, and color are diluted with some of the water and added, with stirring. Sufficient water is then added to produce the
15 required volume.

An aerosol solution is prepared containing the following ingredients:

20 Formulation 5: Aerosol

Ingredient	Quantity (% by weight)
Active ingredient	0.25
Ethanol	25.75
Propellant 22 (Chlorodifluoromethane)	70.00

The active ingredient is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to

-56-

30° C, and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remaining propellant. The valve units are then fitted to the container.

5

Suppositories are prepared as follows:

Formulation 6: Suppositories

Ingredient	Quantity (mg/suppository)
Active ingredient	250
Saturated fatty acid glycerides	2,000

10

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimal necessary heat. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

15

An intravenous formulation is prepared as follows:

Formulation 7: Intravenous Solution

20

Ingredient	Quantity
Active ingredient	50 mg
Isotonic saline	1,000 mL

The solution of the above ingredients is intravenously administered to a patient at a rate of about 1 mL per minute.

25

30

-57-

Formulation 8: Combination Capsule I

Ingredient	Quantity (mg/capsule)
Active ingredient	50
Premarin	1
Avicel pH 101	50
Starch 1500	117.50
Silicon Oil	2
Tween 80	0.50
Cab-O-Sil	0.25

Formulation 9: Combination Capsule II

5

Ingredient	Quantity (mg/capsule)
Active ingredient	50
Norethynodrel	5
Avicel pH 101	82.50
Starch 1500	90
Silicon Oil	2
Tween 80	0.50


Formulation 10: Combination Tablet

Ingredient	Quantity (mg/capsule)
Active ingredient	50
Premarin	1
Corn Starch NF	50
Povidone, K29-32	6
Avicel pH 101	41.50
Avicel pH 102	136.50
Crospovidone XL10	2.50
Magnesium Stearate	0.50
Cab-O-Sil	0.50

-58-

We claim:

1. A compound of Formula I:



wherein:

n is 2 or 3;

R is dimethylamino, diethylamino, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, or 1-hexamethyleneimino;

R^1 is hydrogen, loweralkyl of C_1-C_4 , phenyl, or mono- or disubstituted phenyl wherein each substituent is independently halo, hydroxy, methyl, hydroxy, C_1-C_6 -alkoxy, benzyloxy, C_1-C_6 -alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C_1-C_4 -loweralkyl, or C_1-C_4 -loweralkoxy, C_1-C_5 -alkoxycarbonyloxy, or C_4-C_6 -alkylsulfonyloxy;

R^2 is hydrogen, hydroxy, C_1-C_6 -alkoxy, benzyloxy, C_1-C_6 -alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C_1-C_4 -loweralkyl, or C_1-C_4 -loweralkoxy, C_1-C_5 -alkoxycarbonyloxy, or C_4-C_6 -alkylsulfonyloxy; or a pharmaceutically acceptable acid addition salt thereof.

2. A compound of Claim 1 wherein n is 2.


3. A compound of Claim 1 wherein R is 1-pyrrolidinyl or 1-piperidinyl.

4. A compound of Claim 1 wherein R^1 is methyl.

-59-

5. A compound of Claim 1 wherein R² is hydroxy.
6. A compound of Claim 1 which is 2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-methylpyridine, 2-[4-[2-(1-piperidinyl)ethoxy]phenyl]-3-(4-methoxyphenyl)-6-methylpyridine, 2-[4-[2-(1-piperidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-(4-fluorophenyl)pyridine, 2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3-(4-hydroxyphenyl)-6-phenylpyridine, or 2-[4-[2-(1-pyrrolidinyl)ethoxy]phenyl]-3,6-bis(4-hydroxyphenyl)pyridine, or a pharmaceutically acceptable acid addition salt or solvent thereof.
7. A composition comprising a compound of Claim 1 in combination with a pharmaceutically acceptable carrier, diluent, or excipient, and optionally an effective amount of estrogen or progestin.
8. A method for alleviating the symptoms of post-menopausal syndrome comprising administering to a woman in need of such treatment an effective amount of a compound of Claim 1.
9. A method according to Claim 25 wherein the post-menopausal syndrome pathological condition is osteoporosis, a cardiovascular disease, or estrogen-dependent cancer.
10. A method according to Claim 9 wherein the cardiovascular disease is hyperlipidaemia.
11. A method according to Claim 9 wherein the estrogen-dependent cancer is breast cancer.
12. A method according to Claim 8 comprising the additional step of administering an effective amount of estrogen or progestin.
13. A process for preparing a compound of Formula Ia:

-60-



wherein:


5 n is 2 or 3;

R is dimethylamino, diethylamino, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, or 1-hexamethyleneimino;

10 R^{1a} is hydrogen, loweralkyl of C₁-C₄, phenyl, or mono- or disubstituted phenyl wherein each substituent is independently halo, methyl, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or C₄-C₆-alkylsulfonyloxy;

15 R^{2a} is hydrogen, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzoyloxy, substituted benzoyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or C₄-C₆-alkylsulfonyloxy,

20 which comprises reacting a compound of Formula II:



II


25 with an alkyl chloride of the formula Cl-(CH₂)_n-R, in an inert solvent, in the presence of base, and at a temperature of from 0 to 100°C.

-61-

14. The process of Claim 13 comprising a subsequent step of deprotecting a protected hydroxy substituent.

15. A compound of Formula II:

5



II

wherein:

10 R^{1a} is hydrogen, loweralkyl of C₁-C₄, phenyl, or mono- or disubstituted phenyl wherein each substituent is independently halo, methyl, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy,

benzyloxy, substituted benzyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or 15 C₄-C₆-alkylsulfonyloxy; and

R^{2a} is hydrogen, C₁-C₆-alkoxy, benzyloxy, C₁-C₆-alkanoyloxy, benzyloxy, substituted benzyloxy bearing 1 to 3 substituents each of which is independently halo, C₁-C₄-loweralkyl, or C₁-C₄-loweralkoxy, C₁-C₅-alkoxycarbonyloxy, or 20 C₄-C₆ alkylsulfonyloxy.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/11958

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/44, 31/535; CO7D 213/38, 413/12

US CL :544/124; 546/334; 514/357

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 544/124; 546/334; 514/357

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3,535,329 A (R.G. GRIOT) 20 October 1970, see entire document, especially columns 1 and 2.	1-15
X	US 3,535,330 A (R.G. GRIOT) 20 October 1970, see entire document, especially columns 1-4.	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* "A" document defining the general state of the art which is not considered to be of particular relevance		
* "E" earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* "O" document referring to an oral disclosure, use, exhibition or other means		
* "P" document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search

21 SEPTEMBER 1996

Date of mailing of the international search report

10 OCT 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

R.W. RAMSUER aco

Telephone No. (703) 308-1235